These highlights do not include all the information needed to use DICYCLOMINE HYDROCHLORIDE safely and effectively. See full prescribing information for DICYCLOMINE HYDROCHLORIDE. DICYCLOMINE HYDROCHLORIDE tablets,

Initial U.S. Approval: 1950

----- RECENT MAJOR CHANGES-----Warnings and Precautions, Peripheral and Central

Nervous System (5.3) 07/2012

--- INDICATIONS AND USAGE---Dicyclomine hydrochloride is an antispas and anticholinergic (antimuscarinic) agent indicated for the treatment of functional bowel/ irritable bowel syndrome (1)

-----DOSAGE AND ADMINISTRATION ----

Dosage for dicyclomine hydrochloride must be adjusted to individual patient needs (2). If a dose missed, patients should continue the normal dosing schedule (2).

Oral in adults (2.1):

Kev. U5/2022

Gablets, USP

Dicyclomine

3055-02

Dicyclomine

Hydrochloride

Tablets, USP

Rev. 05/2022

- · Starting dose: 20 mg four times a day. After a week treatment with the starting dose, the dose may be escalated to 40 mg four times a day, unless side effects limit dosage escalation
- Discontinue dicyclomine hydrochloride if efficacy not achieved or side effects require doses less than 80 mg per day after two weeks

---- DOSAGE FORMS AND STRENGTHS----

• Dicyclomine hydrochloride tablets, USP, 20

mg (3) -- CONTRAINDICATIONS

- Infants less than 6 Glaucoma (4) months of age (4) • Obstructive
- Nursing mothers (4) uropathy (4) Obstructive Unstable disease of the
- status in acute hemorrhage (4) tract (4)
- Myasthenia gravis
 Severe ulcerative colitis (4) Reflux esophagitis

----- WARNINGS AND PRECAUTIONS-----

- <u>Cardiovascular conditions</u>: worsening of conditions (5.2)
- Peripheral and central nervous system: heat prostration can occur with drug use (fever and heat stroke due to decreased sweating); drug should be discontinued and supportive measures instituted (5.3)
- Psychosis and delirium have been reported in patients sensitive to anticholinergic drugs (such as elderly patients and/or in patients with mental illness): signs and symptoms resolve within 12 to 24 hours after discontinuation of dicyclomine hydrochloride (5.3)

Myasthenia Gravis: overdose may lead to muscular weakness and paralysis. Dicyclomine hydrochloride should be given to patients with myasthenia gravis only to reduce adverse muscarinic effects of an

- nticholinesterase (5.4) Incomplete intestinal obstruction: diarrhea may be an early symptom especially in patients with ileostomy or colostomy. Treatment with dicyclomine hydrochloride would be inappropriate and possibly fatal
- <u>Salmonella dysenteric patients</u>: due to risk of toxic megacolon (5.6)
- Ulcerative colitis: Dicyclomine hydrochloride should be used with caution in these patients; large doses may suppress intestinal motility or aggravate the serious complications of toxic megacolon (5.7)
- <u>Prostatic hypertrophy</u>: Dicyclomine hydrochloride should be used with caution in se patients; may lead to urinary retention
- Hepatic and renal disease: should be used
- Geriatric: use with caution in elderly who may be more susceptible to dicyclomine hydrochloride's adverse events (5.10)

The most serious adverse reactions include cardiovascular and central nervous system mptoms. The most common adverse reactions > 5% of patients) are dizziness, dry mouth on blurred, nausea, somnolence, asthenia and

To report SUSPECTED ADVERSE REACTIONS, contact TWi Pharn Inc. at 1-844-518-2989 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

---- DRUG INTERACTIONS ----

- Antiglaucoma agents: anticholinergics atagonize antiglaucoma agents and may crease intraocular pressure (7)
- Anticholinergic agents: may affect the gastrointestinal absorption of various drugs; may also increase certain actions or side effects of other anticholinergic drugs (7)
- Antacids: interfere with the absorption of nticholinergic agents (7)

-- USE IN SPECIFIC POPULATIONS -

- Pregnancy: use only if clearly needed (8.1) · Pediatric Use: Safety and effectiveness not
- established (8.4) Hepatic and renal impairment: caution must be taken with patients with significantly impaired hepatic and renal function (8.6)

7.4 Effect of Antacids

8.1 Pregnancy Nursing Mothers

8.4 Pediatric Use

8.5 Geriatric Use

10 OVERDOSAGE

11 DESCRIPTION

8.6 Renal Impairmen

8.7 Hepatic Impairment

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesi

16 HOW SUPPLIED/STORAGE AND

Impairment of Fertility

12.2 Pharmacodynamics

12.3 Pharmacokinetics

See 17 for PATIENT COUNSELING

Effect on Absorption of Other Drugs

7.6 Effect on Gastric Acid Secretic

USE IN SPECIFIC POPULATIONS

FULL PRESCRIBING INFORMATION: CONTENTS*

- 1 INDICATIONS AND USAGE DOSAGE AND ADMINISTRATION
- Oral Dosage and Administration in Adults
- DOSAGE FORMS AND STRENGTHS
- 4 CONTRAINDICATIONS
- WARNINGS AND PRECAUTIONS
- 5.2 Cardiovascular Condition 5.3 Peripheral and Central Nerv
- 5.4 Myasthenia Gravis Intestinal Obstruction
- Toxic Dilatation of Intestine 5.6
- 5.7 Ulcerative Colitis
- Prostatic Hypertrophy
- 5.9 Hepatic and Renal Disease
- 5.10 Geriatric Population

6 ADVERSE REACTIONS

- 6.1 Clinical Trial Experience
- Postmarketing Experience 6.3 Adverse Reactions Reported with

Similar Drugs with Anticholinergic

- Antiglaucoma Agents
- 7.2 Other Drugs with Anticholinergic Activity
- Other Gastrointestinal Motility Drugs
- DRUG INTERACTIONS
 - 17.2 Use in Infants

HANDLING

14 CLINICAL STUDIES

17.3 Use in Nursing Mothers

PATIENT COUNSELING

- 17.4 Peripheral and Central Nervous

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Dicyclomine hydrochloride is indicated for the treatment of patients with functional bowel/irritable

2 DOSAGE AND ADMINISTRATION

Dosage must be adjusted to individual patient needs

2.1 Oral Dosage and Administration in Adults

The recommended initial dose is 20 mg four times a day.

After one week treatment with the initial dose, the dose may be increased to 40 mg four times a day

s side effects limit dosage escalation. If efficacy is not achieved within 2 weeks or side effects require doses below 80 mg per day, the drug

nued. Docume ted safety data are not available for doses about

3 DOSAGE FORMS AND STRENGTHS

Dicyclomine hydrochloride tablets, USP, 20 mg: Light blue to blue round tablet with a few white

4 CONTRAINDICATIONS

Dicyclomine hydrochloride is contraindicated in infants less than 6 months of age [see Use in Specific Populations (8.4)], nursing mothers [see Use in Specific Populations (8.3)], and in patients with:

- unstable cardiovascular status in acute hemorrhage myasthenia gravis [see Warnings and Precautions (5.4)]
- glaucoma [see Adverse Reactions (6.3) and Drug Interactions (7.1)]
- obstructive uropathy [see Warnings and Precautions (5.8)] obstructive disease of the gastrointestinal tract [see Warnings and Precautions (5.5)]
- severe ulcerative colitis [see Warnings and Precautions (5.7)]
- reflux esophagitis

5 WARNINGS AND PRECAUTIONS

5.2 Cardiovascular Conditions

Dicyclomine hydrochloride tablets needs to be used with caution in conditions characterized by tachyarrhythmia such as thyrotoxicosis, congestive heart failure and in cardiac surgery, where they may further accelerate the heart rate. Investigate any tachycardia before administration of dicyclomine hydrochloride tablets. Care is required in patients with coronary heart disease, as ischemia and infarction may be worsened, and in patients with hypertension [see Adverse Reactions (6.3)]

5.3 Peripheral and Central Nervous System

The peripheral effects of dicyclomine hydrochloride tablets are a consequence of their inhibitory The peripheral effects of dicyclomine hydrochloride tablets are a consequence of their limitotry effect on muscarinic receptors of the autonomic nervous system. They include dryness of the mouth with difficulty in swallowing and talking, thirst, reduced bronchial secretions, dilatation of the pupils (mydriasis) with loss of accommodation (cycloplegia) and photophobia, flushing and dryness of the skin, transient bradycardia followed by tachycardia, with palpitations and arrhythmias, and difficulty in micturition, as well as reduction in the tone and motility of the gastrointestinal tract leading to constipation [see Adverse Reactions (6)].

In the presence of high environmental temperature heat prostration can occur with drug use (fever and heat stroke due to decreased sweating). It should also be used cautiously in patients with fever. If ymptoms occur, the drug should be discontinued and supportive measures instituted. Because of the nhibitory effect on muscarinic receptors within the autonomic nervous system, caution should be taken in patients with autonomic neuropathy. Central nervous system (CNS) signs and symptoms include confusional state, disorientation, amnesia, hallucinations, dysarthria, ataxia, coma, euphoria, fatigue, insomnia, agitation and mannerisms, and inappropriate affect.

Psychosis and delirium have been reported in sensitive individuals (such as elderly patients and/or in patients with mental illness) given anticholinergic drugs. These CNS signs and symptoms usually resolve within 12 to 24 hours after discontinuation of the drug.

Dicyclomine hydrochloride may produce drowsiness, dizziness or blurred vision. The patient should be warned not to engage in activities requiring mental alertness, such as operating a motor vehicle or other machinery or performing hazardous work while taking dicyclomine hydrochloride.

With overdosage, a curare-like action may occur (i.e., neuromuscular blockade leading to muscular weakness and possible paralysis). It should not be given to patients with myasthenia gravis except to reduce adverse muscarinic effects of an anticholinesterase [see Contraindications (4)].

Diarrhea may be an early symptom of incomplete intestinal obstruction, especially in patients with ileostomy or colostomy. In this instance, treatment with this drug would be inappropriate and possibly harmful [see Contraindications (4)].

Rarely development of Ogilvie's syndrome (colonic pseudo-obstruction) has been reported. Ogilvie's syndrome is a clinical disorder with signs, symptoms, and radiographic appearance of an acute large bowel obstruction but with no evidence of distal colonic obstruction.

5.6 Toxic Dilatation of Intestine megacolon

Toxic dilatation of intestine and intestinal perforation is possible when anticholinergic agents are administered in patients with Salmonella dysentery.

Caution should be taken in patients with ulcerative colitis. Large doses may suppress intestinal motility to the point of producing a paralytic ileus and the use of this drug may precipitate or aggravate the serious complication of toxic megacolon [see Adverse Reactions (6.3)]. Dicyclomine hydrochloride is contraindicated in patients with severe ulcerative colitis [see Contraindications (4)].

5.8 Prostatic Hypertrophy

Dicyclomine hydrochloride should be used with caution in patients with known or suspected pro enlargement, in whom prostatic enlargement may lead to urinary retention [see Adverse Reactions (6.3)].

5.9 Hepatic and Renal Disease Dicyclomine hydrochloride should be used with caution in patients with known hepatic and renal

Dicyclomine hydrochloride should be used with caution in elderly who may be more susceptible to its

adverse effects.

The pattern of adverse effects seen with dicylomine is mostly related to its pharmacological actions at muscarinic receptors [see Clinical Pharmacology (12)]. They are a consequence of the inhibitory effect on muscarinic receptors within the autonomic nervous system. These effects are dose-related and are usually reversible when treatment is discontinued.

The most serious adverse reactions reported with dicyclomine hydrochloride include cardiovascular and central nervous system symptoms [see Warnings and Precautions (5.2, 5.3)].

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The data described below reflect exposure in controlled clinical trials involving over 100 patients treated for functional bowel/irritable bowel syndrome with dicyclomine hydrochloride tablets at initial doses of 160 mg daily (40 mg four times a day).

In these trials most of the side effects were typically anticholinergic in nature and were reported by 61% of the patients. Table 1 presents adverse reactions (MedDRA 13.0 preferred terms) by decreasing order of frequency in a side-by-side comparison with placebo.

Table 1: Adverse reactions experienced in controlled clinical trials with decreasing order of

MedDRA Preferred Term	Dicyclomine Hydrochloride (40 mg four times a day) %	Placebo %
Dry Mouth	33	5
Dizziness	40	5
Vision blurred	27	2
Nausea	14	6
Somnolence	9	1
Asthenia	7	1
Nervousness	6	2

Nine percent (9%) of patients were discontinued from dicyclomine hydrochloride because of one or more of these side effects (compared with 2% in the placebo group). In 41% of the patients with side effects, side effects disappeared or were tolerated at the 160 mg daily dose without reduction. A dose reduction from 160 mg daily to an average daily dose of 90 mg was required in 46% of the patient nued to experience a favorable clinical response; their side effects either vith side effects who then con

6.2 Postmarketing Experience

The following adverse reactions, presented by system organ class in alphabetical order, have been identified during post approval use of dicyclomine hydrochloride. Because these reactions are reported

ntarily from a population of uncertain size, it is not always possible to reliably estimate their uncy or establish a causal relationship to drug exposure.

- · Cardiac disorders: palpitations, tachyarrhythmia
- · Eye disorders: cycloplegia, mydriasis, vision blurred
- distribution disorders: abdominal distension, abdominal pain, constipation, dry mouth, dyspepsia, nausea, vomiting
- · General disorders and administration site conditions: fatigue, malaise
- · Immune System Disorders: drug hypersensitivity including face oedema, angioedema, anaphylactic
- Nervous system disorders: dizziness, headache, hallucinations insomnia, son
- Psychiatric disorders: As with the other anti-cholinergic drugs, cases of delirium or symptoms
 of delirium such as amnesia (or transient global amnesia), agitation, confusional state, delusion,
 disorientation, hallucination (including visual hallucination) as well as mania, mood altered and
 pseudodementia, have been reported with the use of Dicyclomine. Nervousness and insomnia have also been reported.
- · Respiratory, thoracic and mediastinal disorders: dyspnoea, nasal congestion
- · Skin and subcutaneous tissue disorder: dermatitis allergic, erythema, rash 6.3 Adverse Reactions Reported with Similar Drugs with Anticholinergic/Antispasmodic Action

Central Nervous System: tingling, numbness, dyskinesia, speech disturbance, insomnia

Peripheral Nervous System: With overdosage, a curare-like action may occur (i.e., neuromuscular blockade leading to muscular weakness and possible paralysis)

Ophthalmologic: diplopia, increased ocular tension Dermatologic/Allergic: urticaria, itching, and other dermal manifestations

Genitourinary: urinary hesitancy, urinary retention in patients with prostatic hypertrophy

Respiratory: apnea Other: decreased sweating, sneezing, throat congestion, impotence. With the injectable form, there may be temporary sensation of light-headedness. Some local irritation and focal coagulation necrosis may

occur following the intramuscular injection of dicyclomine hydrochloride.

7.1 Antiglaucoma Agents

Anticholinergics antagonize the effects of antiglaucoma agents. Anticholinergic drugs in the presence of increased intraocular pressure may be hazardous when taken concurrently with agents such as orticosteroids. Use of dicyclomine hydrochloride in patients with glaucoma is not recommended [see Contraindications (4)].

7.2 Other Drugs with Anticholinergic Activity

The following agents may increase certain actions or side effects of anticholinergic drugs including dicyclomine hydrochloride: amantadine, antiarrhythmic agents of Class I (e.g., quinidine), antihistamines, antipsychotic agents (e.g., phenothiazines), benzodiazepines, MAO inhibitors, narcotic analgesics (e.g., meperidine), nitrates and nitrites, sympathomimetic agents, tricyclic antidepressants, and other drugs having anticholinergic activity.

7.3 Other Gastrointestinal Motility Drugs

Interaction with other gastrointestinal motility drugs may antagonize the effects of drugs that alter gastrointestinal motility, such as metoclopramide.

Because antacids may interfere with the absorption of anticholinergic agents including dicyclomine hydrochloride, simultaneous use of these drugs should be avoided.

7.5 Effect on Absorption of Other Drugs

Anticholinergic agents may affect gastrointestinal absorption of various drugs by affecting or gastrointestinal motility, such as slowly dissolving dosage forms of digoxin; increased serum digoxin concentration may result.

7.6 Effect on Gastric Acid Secretion The inhibiting effects of anticholinergic drugs on gastric hydrochloric acid secretion are antagonized by

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Adequate and well-controlled studies have not been conducted with dicyclomine hydrochloride in and two-commended doses of 80 to 160 mg/day. However, epidemiologic studies down an increased risk of structural malformations among babies born to women who took producing dicyclomine hydrochloride tablets at doses up to 40 mg/day during the first trimester of

Reproduction studies have been performed in rats and rabbits at doses up to 33 times the maximum recommended human dose based on 160 mg/day (3 mg/kg) and have revealed no evidence of harm to the fetus due to dieyclomine. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

8.3 Nursing Mothers

nine hydrochloride is contraindicated in women who are breastfeeding. Dicyclomine hydrochloride tablets is excreted in breastmilk. Because of the potential for serious adverse reactions in breastfed infants from dicyclomine hydrochloride, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother [see Use

in Specific Populations (8.4)].

Safety and effectiveness in pediatric patients have not been established.

agents used to treat achlorhydria and those used to test gastric secretion.

Dicyclomine hydrochloride is contraindicated in infants less than 6 months of age [see Contraindications (4)]. There are published cases reporting that the administration of dicyclomine hydrochloride tablets syrup to infants has been followed by serious respiratory symptoms (dyspnea, shortness of breath, breathlessness, respiratory collapse, apnea and asphyxia), seizures, syncope, pulse rate fluctuations, muscular hypotonia, and coma, and death, however; no causal relationship has been established.

Clinical studies of dicyclomine hydrochloride did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the

dosing range in adults, reflecting the greater frequency of decreased hepatic, renal, or cardiac function and of concomitant disease or other drug therapy. Because elderly patients are more likely to have decreased renal function, care should be taken in dose

selection, and it may be useful to monitor renal function

Effects of renal impairment on PK, safety and efficacy of dicyclomine hydrochloride has not been studied. Dicyclomine hydrochloride drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Dicyclomine hydrochloride should be administered with caution in patients with renal impairment.

Effects of renal impairment on PK, safety and efficacy of dicyclomine hydrochloride have not been studied. Dicyclomine hydrochloride should be administered with caution in patients with hepatic

10 OVERDOSAGE In case of an overdose, patients should contact a physician, poison control center (1-800-222-1222), or

The signs and symptoms of overdosage include: headache; nausea; vomiting; blurred vis pupils; hot, dry skin; dizziness; dryness of the mouth; difficulty in swallowing; and CNS stimulatio cluding convulsion. A curare-like action may occur (i.e., neuromuscular blockade leading to muscular weakness and possible paralysis).

One reported event included a 37-year-old who reported numbness on the left side, cold fingertips

tion of 320 mg daily (four 20 mg tablets four times daily). These events resolved after discontinuing

The acute oral LD_{50} of the drug is 625 mg/kg in mice.

The amount of drug in a single dose that is ordinarily associated with symptoms of overdosage or that is likely to be life-threatening, has not been defined. The maximum human oral dose recorded was 600 mg by mouth in a 10-month-old child and approximately 1500 mg in an adult, each of whom survived. In three of the infants who died following administration of dicyclomine hydrochloride tablets [see Warnings and Precautions (5.1)], the blood concentrations of drug were 200, 220, and 505 ng/mL.

It is not known if dicyclomine hydrochloride is dialyzable.


Treatment should consist of gastric lavage, emetics, and activated charcoal. Sedatives (e.g., short-acting barbiturates, benzodiazepines) may be used for management of overt signs of excitement. If indicated, an appropriate parenteral cholinergic agent may be used as an antidote.

11 DESCRIPTION

Dicyclomine hydrochloride, USP, is an antispasmodic and anticholinergic (antimuscarinic) agent available in the following dosage for

Dicyclomine hydrochloride tablets, USP, 20 mg for oral use contain 20 mg dicyclomine hydrochloride, USP. Dicyclomine hydrochloride tablets, USP, 20 mg also contain inactive ingredients: colloidal silicon dioxide, FD&C Blue #1, hypromellose, lactose monohydrate, sodium starch glycolate, and stearic acid

Dicyclomine hydrochloride, USP, is [bicyclohexyl]-1-carboxylic acid, 2-(diethylamino) ethyl ester, hydrochloride, with a molecular formula of $C_{19}H_{35}NO_2$ •HCl and the following structural formula:



Molecular weight: 345.95 Dicyclomine hydrochloride, USP occurs as a fine, white, crystalline, practically odorless powder with a itter taste. It is soluble in water, freely soluble in alcohol and chloroform, and very slightly soluble in

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Dicyclomine relieves smooth muscle spasm of the gastrointestinal tract. Animal studies indicate that this action is achieved via a dual mechanism:

a specific anticholinergic effect (antimuscarinic) at the acetylcholine-receptor sites with approximately 1/8 the milligram potency of atropine (in vitro, guinea pig ileum); and
 a direct effect upon smooth muscle (musculotropic) as evidenced by dieyclomine's antagonism of bradykinin- and histamine-induced spasms of the isolated guinea pig ileum.

Atropine did not affect responses to these two agonists. In vivo studies in cats and dogs showed dicyclomine to be equally potent against acetylcholine (ACh)- or barium chloride (BaCl₂)-induced intestinal spasm while atropine was at least 200 times more potent against effects of ACh than BaCl₂. Tests for mydriatic effects in mice showed that dicyclomine was approximately 1/500 as potent as

atropine; antisialagogue tests in rabbits showed dicyclomine to be 1/300 as potent as atropine

12.3 Pharmacokinetics

Absorption and Distribution

Dicyclomine hydrochloride can inhibit the secretion of saliva and sweat, decrease gastrointestinal etions and motility, cause drowsiness, dilate the pupils, increase heart rate, and depress motor

In man, dicyclomine is rapidly absorbed after oral administration, reaching peak values within 60-90 minutes. Mean volume of distribution for a 20 mg oral dose is approximately 3.65 L/kg suggesting extensive distribution in tissues.

Elimination sm of dicyclomine was not studied. The principal route of excretion is via the urine (79.5% of the dose). Excretion also occurs in the feces, but to a lesser extent (8.4%). Mean half-life of plasma elimination in one study was determined to be approximately 1.8 hours when plasma concentrations were measured for 9 hours after a single dose. In subsequent studies, plasma concentrations were followed for up to 24 hours after a single dose, showing a secondary phase of elimination with a somewhat longer

13 NONCLINICAL TOXICOLOGY ${\bf 13.1\ Carcinogenesis, Mutagenesis, Impairment\ of\ Fertility}$

In studies in rats at doses of up to 100 mg/kg/day, dicyclomine produced no deleterious effects on breeding, conception, or parturition. In controlled clinical trials involving over 100 patients who received drug, 82% of patients treated for functional bowel/irritable bowel syndrome with dicyclomine hydrochloride at initial doses of 160 mg daily (40 mg four times daily) demonstrated a favorable clinical response compared with 55% treated with 150 to 2000.

Long-term animal studies have not been conducted to evaluate the carcinogenic potential of dicyclon

with placebo (p<0.05).

16 HOW SUPPLIED/STORAGE AND HANDLING Dicyclomine hydrochloride tablets, USP, 20 mg is supplied as follows: Light blue to blue round tablet with a few white spots and "T 200" engraved on one side, supplied in bottles of 100, 500 and 1000. To prevent fading, avoid exposure to direct sunlight. Store at 20°-25°C (68°-77°F). [See USP controlled room temperature].

NDC 24979-200-01 NDC 24979-200-02

17 PATIENT COUNSELING INFORMATION 17.2 Use in Infants

Inform parents and caregivers not to administer dicyclomine hydrochloride in infants less than 6 months of age [see Use in Specific Populations (8.4)]. 17.3 Use in Nursing Mothers

Advise lactating women that dicyclomine hydrochloride should not be used while breastfeeding their infants [see Use in Specific Populations (8.3, 8.4)]. 17.4 Peripheral and Central Nervous System In the presence of a high environmental temperature, heat prostration can occur with dicyclomine hydrochloride use (fever and heat stroke due to decreased sweating). If symptoms occur, the drug should be discontinued and a physician contacted. Dicyclomine hydrochloride may produce drowsiness or blurred vision. The patient should be warned not to engage in activities requiring mental alertness, such as operating a motor vehicle or other machinery or to perform hazardous work while taking dicyclomine

as operating a motor vehicle or other machinery or to hydrochloride [see Warnings and Precautions (5.3)]. Manufactured for:

Bora Pharmaceutical Laboratories Inc.

Paramus, NJ 07652

Aptalis Pharma US. Inc.

BENTYL® is a registered trademark owned by Aptalis Pharma Canada Inc., an affiliated company of

Rev. 05/2022

LA-3055-02



Bora Pharmaceutical Laboratories Inc. | Product Name: | Dicyclomine Hydrochloride Tablets Brief 展开尺寸: 412.75X311.15 mm 成品尺寸: 31.75X31.75 mm LA-3055-02 2022-12-05 Item No: 改版 DTP: Job No CSH2022M0025 Roy